arXiv:1210.8337v1 [cond-mat.quant-gas] 31 Oct 2012

Lieb's soliton-like excitations in harmonic trap

G.E. Astrakharchik[1] and L.P. Pitaevskii[2,3]

[1] *Departament de Física i Enginyeria Nuclear, Campus Nord B4-B5, Universitat Politècnica de Catalunya, E-08034 Barcelona, Spain*
[2] *INO-CNR BEC Center and Dipartimento di Fisica, Università di Trento, I-38123 Povo, Trento, Italy*
[3] *Kapitza Institute for Physical Problems RAS, Kosygina 2, 119334 Moscow, Russia.*

PACS `03.75.Lm` – Tunneling, Josephson effect, BoseEinstein condensates in periodic potentials, solitons, vortices, and topological excitations
PACS `05.30.Jp` – Boson systems
PACS `03.75.Kk` – Dynamic properties of condensates; collective and hydrodynamic excitations, superfluid flow

Abstract – We study the solitonic Lieb II branch of excitations in one-dimensional Bose gas in homogeneous and trapped geometry. Using Bethe *ansatz* equations we calculate the "number of particles" and the "effective mass" of a soliton. The frequency of oscillations in a harmonic trap is calculated. It changes continuously from its "soliton-like" value $\omega_h/\sqrt{2}$ in the high density mean field regime to ω_h in the low density Tonks-Girardeau regime with ω_h the frequency of the harmonic trapping. The phase jump of the order parameter is calculated.

Introduction. – Recent development of experimental techniques has opened an exciting possibility to work with ultracold Bose gases in one-dimensional (1D) conditions, for example in a set of elongated optical traps [1] and in magnetic traps created by solid state chips [2]. This development permitted to verify the theoretical predictions in highly controllable experiments. The theoretical investigations of one-dimensional bosons have been started by Marvin Girardeau in Ref. [3], where the case of an infinite repulsion was considered. This case is often called as "Tonks-Girardeau" (TG) limit, although Tonks considered 1D Boltzmann gas [4]. The next important step was made by Lieb and Liniger in Refs. [5, 6] where they obtained an exact solution of the problem of ground state properties and energy spectrum of a one-dimensional Bose gas with the delta-function repulsive interaction (Lieb-Liniger gas). Probably, the most surprising result of the paper [6] is the existence, besides the phonon-like branch of elementary excitations (Lieb I branch), which presence was natural to assume in analogy with 3D case, also of the second branch (Lieb II branch). This branch exists in a finite interval of the momenta $|p|/\rho \leq \pi$ and its energy vanishes, when the coupling constants tends to zero. The meaning of this branch is clarified in the TG-limit, where the Lieb II spectrum coincides with that of an ideal Fermi gas, and corresponds to excitations of holes. The meaning of the second branch in the opposite (mean-field "Bogolyubov") limit of weakly interacting bosons was explained by Kulish, Manakov and Faddeev [7] (see also [8]). They have shown, that the energy-momentum dispersion relation of an excitation of the second branch in this limit coincides with the relation, obtained by Tsuzuki [9] for a soliton, described by the Gross-Pitaevskii equation (GPE). Recently Sato et al. have shown that the spatial profile of the order parameter, defined as a proper matrix element, also reproduces the GPE soliton profile [10]. Hence, Lieb II branch of excitations in the intermediate regime is a result of a quite non-trivial crossover between a topological soliton and an excitation of a fermion-like hole. Investigation of the properties of these unusual objects is, in our opinion, an interesting and important problem. In this paper we will investigate dynamics of Lieb II branch of excitations in a gas confined to 1D harmonic trap. We will assume that the size of the cloud is large in comparison with the healing length. Then one can safely use the Local Density Approximation (LDA) for the dynamics.

Local Density Approximation. – In the LDA, dynamics of an excitation is defined by its dispersion law in a uniform gas. The most convenient description of the dynamics is in terms of energy of the excitation $\varepsilon(V\mu)$ expressed as a function of its velocity V and the chemical potential μ. For a smooth external potential $U(x)$, the LDA energy can be obtained by exchanging the chemical potential by its local value $\mu \to \mu - U(x)$. This means that in the course of the motion of an excitation in the presence of the external field, the energy $\varepsilon(V, \mu - U(x))$ must remain constant [11,12]. Differentiating with respect to time and taking into account that $dx/dt = V$ we obtain $1/V(\partial\varepsilon/\partial V)_\mu dV/dt - (\partial\varepsilon/\partial\mu)_V(\partial U/\partial x) = 0$, or

$$m_{eff}\frac{dV}{dt} = -N_s\left(\frac{\partial U}{\partial x}\right) \tag{1}$$

where the parameters, characterizing the excitations,

$$m_{eff} = \frac{1}{V}\left(\frac{\partial\varepsilon}{\partial V}\right)_\mu \tag{2}$$

$$N_s = -\left(\frac{\partial\varepsilon}{\partial\mu}\right)_V \tag{3}$$

have, correspondingly meaning of the effective mass m_{eff} and the number of atoms N_s in the excitations. For the excitations of the second branch in the Bose gas these quantities are negative, thus $|N_s|$ is the number of atoms expelled at creation of an excitation. Quantities (2) and (3) can be easily calculated in the Bogolyubov regime, where according to [9] the energy of a soliton is $\varepsilon(V,\mu) = 2\hbar(\mu - mV^2)^{3/2}/(3cm^{1/2})$, where $2c$ is the one-dimensional coupling constant (see Eq. (8) below). Correspondingly

$$N_s = -\frac{\hbar}{cm^{1/2}}\left(\mu - mV^2\right)^{1/2}, \tag{4}$$

$$m_{eff} = 2mN_s. \tag{5}$$

Thus, the GPE soliton moves in an arbitrary external field as a particle of the mass $2m$ [12]. If the gas is trapped in a harmonic trap with frequency ω_h, the frequency of oscillations will be $\omega = \omega_h/\sqrt{2}$ [13].

In the opposite TG limit the energy of an second-branch excitation can be presented as $\varepsilon(V,\mu) = \mu - mV^2/2$ and

$$N_s = -1, \tag{6}$$

$$m_{eff} = m \tag{7}$$

corresponding to the "hole-like" nature of the excitation in this limit. In this case $\omega = \omega_h$. In this paper we will calculate the characteristic parameters m_{eff}, N_s and the oscillation frequency ω for intermediate strengths of the interaction.

It is worth to notice, that an excitation, moving in an external field, has a finite lifetime due to decay to several excitations. This effect has been investigated in [14] for the GPE solitons. The probability of the decay is supressed at small enough ω_h. In the following we will not consider this effect.

Lieb's equations. – In the Lieb-Liniger model the Hamiltonian is written as

$$H = \frac{\hbar^2}{2m}\sum_i \frac{d^2}{dx_i^2} + 2c\sum_{i<k}\delta\left(x_i - x_k\right). \tag{8}$$

In the original paper [5] authors used the system of units with $\hbar = 1, m = 1/2$. Calculation of the second branch of the spectrum of elementary excitations is reduced to de solution of a linear integral equation for the function $J(k,q)$

$$2\pi J(k,q) - 2c\int_{-K}^{K}\frac{J(r,q)dr}{c^2 + (r-k)^2} = \pi - 2\tan^{-1}\left(\frac{q-k}{c}\right). \tag{9}$$

The limit of integration K defines the one-dimensional density ρ (and the value of the dimensionless parameter $\gamma = c/\rho$) indirectly, as an integral of the solution of an equation similar (9), but without q-dependent term on the r.h.s. [5]. The dependence $K(\gamma)$ and the inverse one $\gamma(K)$ can be calculated following the methods of [5]. Once such relations are known, the sound velocity u can be calculated according to $u = -2\gamma^2 d(K/\gamma)/d\gamma$. We use matrix methods to solve Eq. (9) and similar integral equations. To do so we discretized the integral which than is written as a (r,q) matrix. The inverse matrix is calculated and is multiplied by the discrete representation of the r.h.s. of Eq. (9).

The knowledge of J permits to calculate the dependence of the energy ϵ on the momentum p in the parametric form (here q is understood as a free parameter):

$$\varepsilon = \mu - q^2 + 2\int_{-K}^{K} J(k,q)k\,dk, \quad p = -q + \int_{-K}^{K} J(k,q)\,dk. \tag{10}$$

The resulting excitation spectrum ε is a function of μ and p, instead of μ and V. It is possible to calculate $V(p,\mu) = (\partial\varepsilon/\partial p)_\mu$ from equations (10). To calculate N_s and m_{eff} in these variables one can use the relations

$$N_s = -\left(\frac{\partial\varepsilon}{\partial\mu}\right)_p + V\left(\frac{\partial V}{\partial\mu}\right)_p\left(\frac{\partial p}{\partial V}\right)_\mu, \tag{11}$$

$$m_{eff} = \left(\frac{\partial\varepsilon}{\partial p}\right)_\mu\left(\frac{\partial p}{\partial V}\right)_\mu. \tag{12}$$

Results and discussion. – Probably, the best way to experimentally verify our predictions is to measure the frequencies of oscillations Ω in a trap in different regimes. Figure 1 shows the dependence of the frequency Ω on the interaction parameter γ. One can see that the frequency continuously increases with increasing γ from its GPE value $\omega_h/\sqrt{2}$ to the ideal Fermi gas value ω_h. The most rapid change takes place at $\gamma \sim 3$. There are different ways of measuring the oscillations frequency. At moderately small values of γ, when a soliton still contains a large number of atoms, one can directly observe its motion, like in the experiments [15]. Instead at $\gamma \sim 1$ the number of atoms in a soliton is small, $|N_s| \sim 1$, one might exploit the confinement induced resonance (CIR) [16] in order to change the value of γ in the course of an experiment. Initially solitons can be created by a phase



.

Fig. 1: (Color online) Solid line, frequency of oscillations Ω in units of the frequency ω_h of the harmonic oscillator as a function of the interaction parameter γ; dashed line, asymptotic value in TG limit; dash-dotted linea, asymptotic value in GP limit; short-dashed line, perturbative solution of integral equations, $\Omega/\omega_h = 1 - 1/\gamma + \ldots$.



.

Fig. 2: (Color online) Solid line, number of particles N_s in a stationary soliton ($V = 0$) as a function of the interaction parameter γ; dashed line TG limit, Eq. (6); dash-dotted line, GP limit, Eq. (4).

imprinting method at small γ. Later the value of γ might be increased by using CIR and soliton oscillations can be excited by a parametric modulation of the trap frequency. One expects to observe a resonance at the frequency of modulation $\Omega/2$. The resonance can be detected by heating of the gas.

The frequency of small oscillations is $\Omega/\omega_h = \sqrt{N_s m/m_{eff}}$. It depends both on the number of particles in the soliton N_s and on the effective mass m_{eff}. These quantities must be taken in the limit of a small velocity $V \to 0$. The dependence of the number of particles in the soliton at rest $N_s(V = 0)$ on the interaction strength is shown in Fig. 2. We find that for small γ (GP regime) $|N_s| \gg 1$ and the soliton is a macroscopic object, however $|N_s|$ becomes of the order of 1 at $\gamma \sim 1$.

At small γ the number of atoms is well described by the GP result, Eq. (4). However, the situation is different for "fast" solitons with a vanishing momentum $p \to 0$, i.e. with the velocity approaching the speed of sound $V \to u$. According to Eq. (4) the number of atoms in a soliton tends to zero, $|N_s| \propto (u - V)^{1/2}$. However, the calculations show that N_s tends to a finite value at $V \to u$. It is not difficult to prove, that this is only possible, if the dispersion law of the soliton has the following expansion at $p \to 0$

$$\varepsilon(p) \approx up + \frac{p^2}{2m^*}, \tag{13}$$

with $m^* = |m_{eff}(p = 0)|$. This relation is quite nontrivial, because presence of the p^2 contradicts the GPE. Indeed, according to GPE $\varepsilon - up \propto p^{5/3}$. The existence of the p^2 term was established by Periera et al. [17] and Imambekov, Schmidt, and Glazman (see Eqs. (50)-(51) in [18]). Such a term exists both for upper and lower branches of elementary excitations. The effective mass is the same in the absolute value for two branches. Simple calculation permits to present the result of [18] as

$$|m_{eff}(p = 0)|^{-1} = \frac{3}{4}\sqrt{\frac{u}{\pi\hbar m\rho}}\left(1 + \frac{\rho}{3u^2}\frac{d(u^2/\rho)}{d\rho}\right) . \tag{14}$$

In the GPE regime $\gamma \ll 1$, velocity of sound $u \propto \rho^{1/2}$ and the second term disappears. Taking to account that the presence of the p^2 term in dispersion does not violate the relation $N_s = m_{eff}/2$, one finds for $\gamma \to 0$ [18]

$$|N_S(p = 0)| = \frac{2\sqrt{\pi}}{3}\gamma^{-1/4} = 1.18\gamma^{-1/4} . \tag{15}$$

In Fig. 3 we test the obtained result by showing the dependence of $|N_S(p = 0)|\gamma^{1/4}$ on γ. One can see a good agreement for the coefficient in Eq. (15) in GP limit.

In Galilean-invariant systems with a non-diagonal long range order, there exists an important relation between the momentum of a solition, the number of atoms N_s and a topological characteristic of the soliton, namely the phase jump φ_s. This relation must be valid both in a superfluid Bose system [19] and in a superfluid Fermi gas [20]. For a Bose gas the relation can be written as

$$\hbar\rho\varphi_s = p - mN_sV . \tag{16}$$

The phase jump as a function of the momentum p for different values of γ is presented in Fig. 4.

Notice that in the TG limit large γ the jump $\varphi_s = \pi$ and does not depend on V. We could not find a simple interpretation of this result.

The derivation of Eq. (16) implies an existence of a well-defined complex order parameter. The validity of the



.

Fig. 3: Solid line, number of particles in the soliton N_s for zero momentum $p = 0$ (or soliton velocity $V = u$) scaled with $\gamma^1/4$ as a function of γ. Dashed line, constant value of 1.18 as expected from Eq. (15) in GP regime.

equation in a 1D system at not small values of γ, where the order parameter is destroyed due to the long-distance phase fluctuations, is an interesting open question. In this connection it would be quite interesting to compare these results with the results of direct numerical simulation of the quantum dynamics of solitons at moderate γ, analogous to presented in [21] and with the exact calculations of the profile of the order parameter by the method of [10].

Conclusions. – To conclude, by using the exact Lieb-Liniger theory we investigate physical characteristics of the motion of soliton-like Lieb II branch of excitations. The frequency of oscillations, number of atoms in the soliton and phase jump are calculated. These quantities experience a continuous change in the crossover between Gross-Pitaevskii and Tonks-Girardeau limits. The experimental possibility of the verification of the calculations is discussed.

* * *

Authors thank L. D. Faddeev and L. I. Glazman for fruitful discussions. G.E.A. acknowledges support from Ministerio de Educación, Cultura y Deporte (MEC) (Spain) through the Ramón y Cajal fellowship program and (Spain) Grant No. FIS2011-25275. L.P.P. acknowledges support from the ERC through the QGBE grant.

REFERENCES

[1] Moritz H., Stöferle T., Köhl M. and Esslinger T., *Phys. Rev. Lett.*, **91** (2003) 250402.



.

Fig. 4: (Color online) Phase jump φ_s as a function of soliton momentum p for different values of the interaction strength γ. Upper curves correspond to larger values of γ, lower curves to smaller. The phase jump is constant $\phi_s = \pi$ in the TG limit $\gamma \to \infty$.

[2] Esteve J., Trebbia J.-B., Schumm T., Aspect A., Westbrook C. I. and Bouchoule I., *Phys. Rev. Lett.*, **96** (2006) 130403.

[3] Girardeau M., *J. Math. Phys. (NY)*, **1** (1960) 516.

[4] Tonks L., *Phys. Rev.*, **50** (1936) 955.

[5] Lieb E. H. and Liniger W., *Phys. Rev.*, **130** (1963) 1605.

[6] Lieb E. H., *Phys. Rev.*, **130** (1963) 1616.

[7] Kulish P. P, Manakov S. V. and Faddeev L. D., *Theoreticheskaya i Matematicheskaya Fizika*, **28** (1976) 38; [*Theoretical and Mathematical Physics*, **28** (1976) 615].

[8] Ishikawa M. and Takayama H., *J. Phys. Soc. Jap.*, **49** (1980) 1242.

[9] Tsuzuki T., *J. Low Temp. Phys.*, **4** (1971) 441.

[10] Sato J., Kanamoto R., Kaminishi E. and Deguchi T., *arXiv:1204.3960v1*, (2012) .

[11] Fedichev P. O., Muryshev A. E. and Shlyapnikov G. V., *Phys. Rev. A*, **60** (1999) 3220.

[12] Konotop V. V. and Pitaevskii L., *Phys. Rev. Lett.*, **93** (2004) 240403.

[13] Busch Th. and Anglin J. R., *Phys. Rev. Lett.*, **84** (2000) 2298.

[14] Wadkin-Snaith D. C. and Gangardt D. M., *Phys. Rev. Lett.*, **108** (2012) 085301.

[15] Burger S., Bongs K., Dettmer S., Ertmer W., Sengstock K., Sanpera A., Shlyapnikov G. V. and Lewenstein M., *Phys. Rev. Lett.*, **83** (1999) 3577.

[16] Olshanii M., *Phys. Rev. Lett.*, **81** (1998) 938.

[17] Pereira R. G., Sirker, J., Caux J.-S., Hagemans R., Maillet J. M., White S. R. and Affleck I., *Phys. Rev. Lett.*, **96** (2006) 257202.

[18] Imambekov A., Schmidt T. L. and Glazman L. I., *Rev. Mod. Phys.*, **84** (2012) 1253.

[19] Shevchenko S., *Sov. J. Low Temp. Phys.*, **14** (1988) 553.

[20] Scott R. G., Dalfovo F., Pitaevskii L. P. and

Stringari S., *Phys. Rev. Lett.*, **106** (2011) 185301.
[21] Martin A. D. and Ruostekoski J., *Phys. Rev. Lett.*, **104** (2010) 194102.